

Monograph on
**Endemism in the
Highlands and Escarpments
of Angola and Namibia**



Angola Cave-Chat *Xenocopsychus ansorgei*
Photo: M Mills

Editors:

John M Mendelsohn
Brian J Huntley
Pedro Vaz Pinto

Published with support and funding from:

Ongava Research Centre (ORC)
Namibian Chamber of Environment (NCE)
Centro de Investigação em Biodiversidade
e Recursos Genéticos (CIBIO)
B2Gold Namibia
TotalEnergies

Language editor: Carole Roberts
Design and layout: Alice Jarvis

NE Namibian Journal
JE of Environment

2023: Volume 8 www.nje.org.na

ISSN: 2026-8327 (online)

» [DOWNLOAD THE MONOGRAPH](#)

CONTENTS

Huntley BJ, Mendelsohn JM & Vaz Pinto P Preface to endemism on the highlands and escarpments of Angola and Namibia	i–iii
Huntley BJ, Mendelsohn JM & Vaz Pinto P The biological importance of the highlands of Angola and Namibia: Synopsis and conclusions.....	v–xiii

Geography of the highlands and escarpments

Jarvis AM The highlands and escarpments of Angola and Namibia: orientation maps.....	1–6
Mendelsohn JM & Huntley BJ Introducing the highlands and escarpments of Angola and Namibia	7–22
Miller RM Geology and landscape evolution of the highlands and escarpments of western Angola and Namibia.....	23–28
Huntley BJ Biomes and ecoregions of the highlands and escarpments of Angola and Namibia	29–41
Mendelsohn JM & Gomes AL The human environment in the highlands and escarpments of Angola and Namibia	43–51
Vaz Pinto P, Russo V & Veríssimo L The highlands in Angolan conservation areas	53–62

Diversity and endemism

Craven P & Kolberg H An overview of plant endemism on the highlands of Namibia	63–76
Goyder DJ, Gomes AL, Gonçalves FMP, Luís JC & Darbyshire I A botanical assessment of Mt Namba, Cuanza-Sul, Angola: an isolated mountain towards the northwestern limits of the Great Escarpment of southern Africa.....	77–92
Meller P, Lages F, Finckh M, Gomes A & Goyder D Diversity and endemism of geoxylic plants on the Angolan Planalto.....	93–109
Bruyns PV, Hanáček P & Klak C Diversity and endemism in the species-rich Ceropegieae (Apocynaceae) and <i>Euphorbia</i> in the highlands and escarpments of Angola and Namibia	111–134
Dexter KG, Swanepoel W, Loiseau O, Darbyshire I, Nanyeni L, Gonçalves FM, Chase F & Manzitto-Tripp EA High endemism of the genus <i>Petalidium</i> (Acanthaceae) in the highlands and escarpments of Angola and Namibia	135–147
Weeks A & Swanepoel W <i>Commiphora</i> of the highlands and escarpments of Angola and Namibia	149–159
Lautenschläger T, Aime MC, Clausnitzer V, Langer L, Meller P, Müller F, Nuss M, Teutloff N & Ernst R Green gem of the Northern Escarpment: biodiversity and endemism of the Serra do Pingano Forest Ecosystem.....	161–172
Kipping J, Clausnitzer V & Dijkstra K-DB The highlands and escarpment of Angola as an endemism hotspot for African dragonflies and damselflies (Insecta: Odonata).....	173–186
Gunter F, Jürgens N & Henschel JR Observations on the diversity of termites in Angola and Namibia.....	187–192
Mansell MW The Neuroptera of the highlands and escarpments of Angola and Namibia	193–196
Gomez K, Hawkes PG & Fisher BL Ant endemism in the highlands and escarpments of Angola and Namibia (Hymenoptera, Formicidae)	197–203
Gardiner AJ & Williams MC The endemic butterflies of Angola and Namibia and their evolutionary implications.....	205–230
Prendini L & Bird TL Endemism of Arachnida (Amblypygi, Scorpiones and Solifugae) in the highlands and escarpments of Angola and Namibia: current knowledge and future directions.....	231–244
Becker FS, Baptista NL, Vaz Pinto P, Ernst R & Conradie W The amphibians of the highlands and escarpments of Angola and Namibia.....	245–257
Bauer AM, Ceriáco LMP, Marques MP & Becker FS Highland reptiles of Angola and Namibia	259–276
Conradie W, Lobón-Rovira J, Becker FS, Schmitz A & Vaz Pinto P Flat gecko (<i>Afroedura</i>) diversity, endemism and speciation in the highlands and escarpments of Angola and Namibia.....	277–281
Skelton PH Fishes of the highlands and escarpments of Angola and Namibia.....	283–292
Mills MSL & Melo M Birds of the highlands and escarpments of Angola and Namibia: ornithological significance, avifaunal patterns and questions requiring further study	293–309
Palmeirim AF, Monadjem A, Vaz Pinto P, Taylor P, Svensson MS & Beja P Mammal endemism in the highlands and escarpments of Angola and Namibia.....	311–322
De Matos D, Zastrow J, Val A & Mendelsohn JM Caves and their fauna in the highlands and escarpments of Angola and Namibia	323–330

The Neuroptera of the highlands and escarpments of Angola and Namibia

MW Mansell

URL: <https://www.nje.org.na/index.php/nje/article/view/volume8-mansell>

Published online: 15th December 2023

Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa; mansel@mweb.co.za

ABSTRACT

Of the 17 families of Neuroptera (lacewings) worldwide, ten families represented by 148 species have been recorded from Namibia, and five families with 36 species from Angola. In Namibia, 17 endemic species are known, of which two are known from high-elevation escarpment localities. However, very few surveys dedicated specifically to Neuroptera have been carried out in Namibia and none in Angola, so it is therefore not possible to assess the importance of the highlands and escarpments to Neuroptera. Recommendations are made for further surveys and research.

Keywords: Angola, escarpments, highlands, lacewings, Namibia, Neuroptera

INTRODUCTION

The Neuroptera, generally referred to as ‘lacewings’, is one of 28 orders that comprise the animal class Insecta – the insects. The Neuroptera includes 17 families distributed worldwide, with exceptionally diverse faunas in southern Africa, especially in South Africa and Namibia which are both endowed with a rich diversity of families and species, reflecting a recent evolutionary diversification in the southern part of Africa. Ten families of Neuroptera have been recorded from Namibia and five from Angola.

GENERAL BIOLOGY OF NEUROPTERA

The name ‘Neuroptera’ is derived from the Greek *neuron* which means nerve or sinew and *ptera* which means wings, thus the name means ‘nerve-winged insects’ referring to the characteristic lace-like pattern of veins in the wings. The adults can be easily recognised by the two pairs of similar diaphanous wings that are held in a roof-like manner over the body when the insects rest with the wings folded. The antennae are also definitive, ranging from long, multi-segmented and filiform, to short and stout, to long and clubbed; features that clearly separate them from the similar-looking dragonflies and damselflies.

Adult lacewings are either predacious with biting and chewing mouthparts, or pollen and nectar feeders having mouthparts adapted for pollen and nectar collection, especially the Nemopteridae.

Lacewing larvae, by contrast, are all obligate predators, with piercing and sucking mouthparts, comprising a complementary combination between the mandibles and the lacinia of the maxillae, the latter slotting into a groove on the ventral surface of the mandibles to form a tubular canal (Figure 1) through which saliva, containing digestive enzymes

can pass. The resulting liquid food can then be drawn into the alimentary canal. This feature is a highly evolved, albeit ancient, evolutionary advance which is unique to the Neuroptera, and provides the autapomorphy (i.e., specialised character or trait that is unique to a monophyletic taxonomic group) that defines the lacewings as a monophyletic evolutionary lineage. When a lacewing larva detects prey, it uses its mandibles to seize the prey, and then injects proteolytic enzymes into the prey’s body. Once the internal organs are liquified, the resultant fluid is sucked into the alimentary canal of the larva, where further digestion takes place. This digestive process is highly efficient and the larvae do not excrete solid matter during the entire larval life, which can last from a few weeks to several years, depending on the species. Accumulated solid matter is voided as a single meconial pellet when the larva metamorphoses into an adult and emerges from the cocoon. This digestive system has also enabled the Malpighian tubules to depart from their normal excretory



Figure 1: Lacewing larvae are all obligate predators with piercing and sucking mouthparts, comprising a complementary combination between the mandibles and the lacinia of the maxillae, the latter slotting into a groove on the ventral surface of the mandibles to form a tubular canal. Ventral view of head of *Palpares inclemens*.

function and to evolve into silk-producing glands, another unique autapomorphic feature of the Neuroptera. All lacewing larvae spin a spherical silken cocoon in which to pupate. The cocoon protects the metamorphosing larva from parasites and predators, by affording camouflage and reducing dehydration.

While evolutionary advances such as these often lead to very limited possibilities of further biological advances, or even evolutionary cul-de-sacs, lacewings have used them to adapt to a variety of ecological niches, leading to further evolutionary adaptations and advances. They consequently provide numerous examples of habitat, biological and behavioural diversifications, which have ensured their survival by diversification through specialisation.

Habitats and life strategies

The morphological and physiological adaptations have enabled lacewings, especially the larvae, to evolve into a diversity of habitats and life strategies, including: aquatic, where larvae feed on freshwater sponges; semi-aquatic on the borders of streams; as inquilines in the nests of ants and termites; as parasites in spiders' nests; as arboreals, living on branches and foliage; and living in tree holes, caves, rock overhangs and in sand.

Significance of lacewings in ecosystems

Insects are arguably the most significant component of the biodiversity and functioning of ecosystems. They render numerous ecosystem services, ranging from pollination, nutrient recycling through biodegradation of organic matter, to population control of other invertebrates and plants. Most plants are insect-pollinated and all depend on recycled nutrients. Insects are also vital components of the food chains of other animals, especially birds, fishes and other vertebrate animals. Adults of many Neuroptera are pollinators of indigenous plants, with some having co-evolved with certain plants. The larvae are all obligate predators and limit populations of other insects and invertebrates. They are also important indicators of ecosystem health and are a significant component of the biodiversity of many ecosystems, including arid areas.

NEUROPTERA IN NAMIBIA AND ANGOLA

Namibia

Of the 17 worldwide families of Neuroptera, 10 have been recorded from Namibia, with the largest representation being the families usually known as antlions (Myrmeleontidae), thread- and ribbon-wing lacewings (Nemopteridae) and green lacewings (Chrysopidae). Other families represented in Namibia include: silky or moth lacewings (Psychopsidae; three species), owlflies (Ascalaphidae; ten species), dusty lacewings (Coniopterygidae; eight species), sponge

flies (Sisyridae; one species), beaded lacewings (Berothidae; three species), raptor lacewings (Rhachiberothidae; three species) and pleasing lacewings (Dilaridae; one species). All of these families are also known from South Africa, and other southern African countries.

The ten families of Neuroptera in Namibia are represented by 148 species. Of these, 50 species – more than one third – were originally described from Namibia and 17 are endemic to Namibia. The remaining 98 taxa are more widespread and are also known from neighbouring countries, especially South Africa, with a few also occurring in Botswana and Angola.

Figure 2 illustrates some of the species which are endemic to Namibia. Of the 17 Namibian endemics, only two, *Tjederia brevicornis* (Nemopteridae, Crocinae) (Mansell 1981) and an undescribed species of *Tricholeon* (Myrmeleontidae) are specifically known from high-lying escarpment localities. *Tjederia brevicornis* occurs in the mountains around Windhoek, while the *Tricholeon* species is the only endemic lacewing known from the Brandberg (Mansell & Aspöck 1990).

Namibian surveys

Until recently, very few surveys dedicated specifically to Neuroptera were carried out in Namibia:

- The British Museum expedition of southern Africa during the 1950s included sites in Namibia. Species from this expedition were described by the great Swedish neuropterist Bo Karl Herman Tjeder, in a series of fundamental works entitled “South African Animal Life” (Tjeder 1960, 1961, 1966, 1967).
- The current author, M Mansell, did surveys during 1970–1990, that were specifically focused on the study of larvae of thread-winged lacewings (Nemopteridae, Crocinae). A wide area was covered, which included mountainous areas in and near Windhoek and the Khomas Hochland (Mansell 1981).
- In 1988, the Third International Symposium on Neuropterology was held in South Africa and included a post-symposium excursion to Namibia during which some of the world's leading exponents of Neuroptera conducted limited collections (Mansell & Aspöck 1990).

Prior to these collections, F Gaerdes and S Braine collected many species of Neuroptera that contributed to the discourse, and provided several specimens upon which new species descriptions were based. More recently, surveys have been carried out by R Becker and A Moller through their Lacewing

Monitoring Project, and by the Skeleton Coast–Iona Transfrontier Park Technology for Conservation (SCIONA) Project of the Namibia University of Science and Technology.

The survey of the Brandberg by E Marais and A Kirk-Spriggs in 1998, is the only study that has specifically inventoried any inselberg or high-lying areas in Namibia. The Neuroptera were particularly well represented in the insect samples from these collections and were comprehensively documented




Figure 2: A selection of Namibian Neuroptera species: a) *Isonemurus longipalpis*, a rare Namibian endemic (photo: M Mansell); b) *Crambomorphus namibicus*, a beautiful Namibian endemic, habitus (photo: M Mansell); c) *Crambomorphus namibicus*, profile, habitus (photo: M Mansell); and d) *Crambomorphus namibicus* larva, rarely encountered (photo: R Oberprieler).

by Mansell (2000). It included a complete list of the Neuroptera known from Namibia at that time, and specifically listed 40 species in seven families from the Brandberg. However, all of the species recorded from the Brandberg occur elsewhere in Namibia, with the exception of a solitary adult specimen of an undescribed species of *Tricholeon*, (Myrmeleontidae) which may prove to be a Brandberg endemic (Mansell 2000).

Angola

There has never been any attempt to survey Angola for Neuroptera, and all of the known species are the result of fortuitous bycatch. The neuropteran fauna of Angola is the most poorly known on the African continent and only five families have been recorded: Myrmeleontidae, Nemopteridae, Psychopsidae, Ascalaphidae and Rhachiberothidae. Most of these are known from very few records, three or fewer, in the literature. Of the 36 species known, 10 (almost one third) were originally described from specimens collected in Angola, and most are still only known by the unique type specimens. Nothing is, unfortunately, known about the Neuroptera of the escarpment areas in Angola.

DEFICIENCIES IN KNOWLEDGE, PRIORITIES FOR RESEARCH AND CONSERVATION

The main deficiency in terms of knowledge of Neuroptera in the highlands and escarpments of Angola and Namibia (HEAN) is obviously the lack of records from the highlands and escarpments of both countries. Without such records no assessment of the importance of the highlands and escarpments is possible. Thus dedicated surveys of specific areas are the first priority for research. Such surveys should target both adult specimens and larvae to gain a better understanding of their biology in particular areas. These could be conducted at the same time as surveys of other organisms, especially insects and other invertebrates. As there is essentially no knowledge of the Neuroptera in Angola in general, a wider priority should be to address this deficiency as well.

The author has expertise on the methods required to collect specimens and material for DNA analyses and could also provide taxonomic and biological information support for future surveys. Some databases are already available to provide some data, especially from Namibia. Regarding Angola, it would also be advisable to determine whether there are any specimens held in collections there, for example, in the museum in Luanda or other repositories.

For Namibian lacewings, an immediate priority would be to locate the larvae of the endemic species of *Tricholeon* in the Brandberg, and to determine

through larval collections whether this species also occurs elsewhere. As we already know what the larval requirements of other species in the genus are, this would be relatively easy to achieve in the short term.

ACKNOWLEDGEMENTS

The author is grateful to Rolf Becker, Alma Moller and Vera de Cauwer for enhancing knowledge by facilitating further collections under the auspices of the SCIONA and Namibian Neuroptera projects.

REFERENCES

- Mansell MW (1981) The Crocinae of southern Africa (Neuroptera: Nemopteridae). 3. The genus *Tjederia* Mansell, with keys to the southern African Crocinae. *Journal of the Entomological Society of Southern Africa* 44(2): 245–257.
- Mansell MW (2000) Neuroptera (Insecta). In: Kirk-Spriggs AH, Marais E (eds) Däures: biodiversity of the Brandberg Massif, Namibia. *Cimbebasia Memoir* 9: 163–176. Windhoek.
- Mansell MW, Aspöck H (1990) Post-symposium neuropterological excursions. In: Mansell MW, Aspöck H (eds) *Advances in neuropterology. Proceedings of the Third International Symposium on Neuropterology. Meeting: 3-4 February 1988, Berg en Dal, Kruger National Park, South Africa. South African Department of Agricultural Development, Pretoria.* vi + 298 pp.
- Tjeder B (1960) Neuroptera-Planipennia. The lace-wings of southern Africa. 3. Family Psychopsidae. In: Hanström B, Brinck P, Rudebec G (eds) *South African animal life.* 7: 164–209. Swedish Natural Science Research Council, Stockholm.
- Tjeder B (1961) Neuroptera-Planipennia. The lace-wings of southern Africa. 4. Family Hemerobiidae. In: Hanström B, Brinck P, Rudebec G (eds) *South African animal life.* 8: 296–408. Swedish Natural Science Research Council, Stockholm.
- Tjeder B (1966) Neuroptera-Planipennia. The lace-wings of southern Africa. 5. Family Chrysopidae. In: Hanström B, Brinck P, Rudebec G (eds) *South African animal life.* 12: 228–534. Swedish Natural Science Research Council, Stockholm.
- Tjeder B (1967) Neuroptera-Planipennia. The lace-wings of southern Africa. 6. Family Nemopteridae. In: Hanström B, Brinck P, Rudebec G (eds) *South African animal life.* 13: 290–501. Swedish Natural Science Research Council, Stockholm.